

Windbreak Interpretations

Farmstead windbreaks protect livestock, buildings, and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, increase crop yields, and provide food and cover for wildlife.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth based on the soil type and climate.

Information in this subsection, which includes windbreak suitability groups and a windbreak and environmental plantings table, can be used as a guide in planning windbreaks and screens.

This subsection includes:

- **(a) Trees and Shrubs for Windbreak and Environmental Plantings**

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
1C: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
1D2: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
1E2: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
1F: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
2F: Clarksville-----	---	---	---	---	other trees
Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
3C: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
3D2: Menfro, ERODED-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
3E2: Menfro, ERODED-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
4D2: Menfro, ERODED-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
Bucklick, ERODED-----	---	---	---	---	---
4E: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
Bucklick-----	---	---	---	---	---
5C: Minnith-----	---	---	---	---	---
7E2: Menfro, ERODED-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
Caneyville, ERODED-----	---	---	---	---	---
8D2: Minnith, ERODED-----	---	---	---	---	---
Lily, ERODED-----	---	---	---	---	---

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
9C: Hildebrecht-----	American plum; common lilac; fragrant sumac	Amur maple; gray dogwood; Washington hawthorn	Austrian pine; common hackberry; eastern redcedar; honeylocust; unknown; Virginia pine	---	---
9D: Hildebrecht-----	American plum; common lilac; fragrant sumac	Amur maple; gray dogwood; Washington hawthorn	Austrian pine; common hackberry; eastern redcedar; honeylocust; unknown; Virginia pine	---	---
10F: Menfro-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
Caneyville-----	---	---	---	---	---
Rubble-----	---	---	---	---	---
11E: Goss-----	fragrant sumac	American plum; gray dogwood; southern arrowwood	eastern redbud; eastern redcedar; Washington hawthorn	green ash; northern red oak; tuliptree; white fir	eastern white pine
12E: Lily-----	---	---	---	---	---
Minnith-----	---	---	---	---	---
15E: Gasconade-----	---	---	---	---	other trees
Rock outcrop-----	---	---	---	---	---

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
18C: Weller-----	---	---	---	---	---
19C: Weingarten-----	---	---	---	---	---
19D: Weingarten-----	---	---	---	---	---
20: Wilbur, FREQUENTLY FLOODED-----	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak
21: Haymond, FREQUENTLY FLOODED-----	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak
24A: Elsah, FREQUENTLY FLOODED-----	---	---	---	---	---
25A: Midco, FREQUENTLY FLOODED-----	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak
Riverwash-----	---	---	---	---	---
26A: Auxvasse, RARELY FLOODED	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
27A: Ashton, RARELY FLOODED--	---	---	---	---	---
28A: Freeburg, RARELY FLOODED	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak
29: Kickapoo, FREQUENTLY FLOODED-----	---	---	---	---	---
32E: Gasconade-----	---	---	---	---	---
Caneyville-----	---	---	---	---	---
50: Darwin, RARELY FLOODED--	buttonbush	possumhaw	eastern arborvitae; eastern redcedar; nannyberry	baldcypress; common hackberry; pin oak	eastern cottonwood
52: Parkville, RARELY FLOODED-----	American plum; blackhaw	possumhaw	eastern arborvitae; eastern redcedar; nannyberry; Washington hawthorn; white spruce	bur oak; common hackberry; green ash	---
53: Leta, RARELY FLOODED----	---	---	---	---	---
54: Waldron, RARELY FLOODED-	---	---	---	---	---
55: Haynie, RARELY FLOODED--	---	---	---	---	---

Perry County, Missouri
Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
58: Dupo, RARELY FLOODED----	American plum; fragrant sumac	blackhaw; gray dogwood	eastern redcedar; nannyberry; Washington hawthorn	baldcypress; green ash; sweetgum	eastern white pine; pin oak
62A: Haynie, RARELY FLOODED--	---	---	---	---	---
Waldron, RARELY FLOODED-	---	---	---	---	---
65: Orthents-----	---	---	---	---	---
Water-----	---	---	---	---	---
66: Pits-----	---	---	---	---	---
AED: Orthent-----	---	---	---	---	---
W: Water-----	---	---	---	---	---